arXiv:0904.2916v12 [math.AG] 12 Jul 2010

ON JETS, EXTENSIONS AND CHARACTERISTIC CLASSES I

HELGE MAAKESTAD

Abstract. In this paper we give general definitions of non-commutative jets in the local and global situation using square zero extensions and derivations. We study the functors $\mathrm{Exan}_k(A, I)$ where A is any k-algebra and I is any left and right A-module and use this to relate affine non-commutative jets to liftings of modules. We also study the Kodaira-Spencer class $\mathrm{KS}(\mathcal{L})$ and relate it to the Atiyah class.

Contents

1. Introduction 1
2. Jets, liftings and small extensions 1
3. Atiyah classes and Kodaira-Spencer classes 10
References 15

1. Introduction

In this paper we give general definitions of non-commutative jets in the local and global situation using square zero extensions and derivations. We study the functors $\mathrm{Exan}_k(A, I)$ where A is any k-algebra and I is any left and right A-module and use this to relate affine non-commutative jets to liftings of modules. In the final section of the paper we define and prove basic properties of the Kodaira-Spencer class $\mathrm{KS}(\mathcal{L})$ and relate it to the Atiyah class.

2. Jets, liftings and small extensions

We give an elementary discussion of structural properties of square zero extensions of arbitrary associative unital k-algebras. We introduce for any k-algebra A and any left and right A-module I the set $\mathrm{Exan}_k(A, I)$ of isomorphism classes of square zero extensions of A by I and show it is a left and right module over the center $C(A)$ of A. This structure generalize the structure as left $C(A)$-module introduced in [3]. We also give an explicit construction of $\mathrm{Exan}_k(A, I)$ in terms of cocycles. Finally we give a direct construction of non-commutative jets and generalized Atiyah sequences using derivations and square zero extensions.

Let in the following k be a fixed base field and let

$$0 \to I \to B \to A \to 0$$

Date: Spring 2009.

1991 *Mathematics Subject Classification.* 14F10, 14F40.

Key words and phrases. Atiyah sequence, jet bundle, Lie-Rinehart algebra, Chern class, Atiyah class, Kodaira-Spencer class, square zero extension, lifting.

Partially supported by a scholarship from NAV, www.nav.no.

be an exact sequence of associative unital k-algebras with $i(I)^2 = 0$. Let $i : I \to B$ and $p : B \to A$ denote the morphisms. Assume s is a map of k-vector spaces with the following properties:

$$s(1) = 1$$

and

$$p \circ s = \mathrm{id}\,.$$

Such a section always exist since B and A are vector spaces over the field k. Note: s gives the ideal I a left and right A-action.

Lemma 2.1. *There is an isomorphism*

$$B \cong I \oplus A$$

of k-vector spaces.

Proof. Define the following maps of vector spaces: $\phi : B \to I \oplus A$ by $\phi(x) = (x - sp(x), p(x))$ and $\psi : I \oplus A \to B$ by $\psi(u, x) = u + s(x)$. It follows $\psi \circ \phi = \mathrm{id}$ and $\phi \circ \psi = \mathrm{id}$ and the claim of the Proposition follows. □

Define the following element:

$$\tilde{C} : A \times A \to I$$

by

$$\tilde{C}(x \times y) = s(x)s(y) - s(xy).$$

It follows that $\tilde{C} = 0$ if and only if s is a ring homomorphism.

Lemma 2.2. *The map $\tilde{C}$ gives rise to an element $C \in \mathrm{Hom}_k(A \otimes_k A, I)$.*

Proof. We easily see that $\tilde{C}(x+y, z) = \tilde{C}(x, z)+\tilde{C}(y, x)$ and $\tilde{C}(x, y+z) = \tilde{C}(x, y)+\tilde{C}(x, z)$ for all $x, y, z \in A$. Moreover for any $a \in k$ it follows

$$\tilde{C}(ax, y) = \tilde{C}(x, ay) = a\tilde{C}(x, y).$$

Hence we get a well defined element $C \in \mathrm{Hom}_k(A \otimes_k A, I)$ as claimed. □

Define the following product on $I \oplus A$:

$$(u, x) \times (v, y) =(uy + xv + C(x, y), xy). \tag{2.2.1}$$

We let $I \oplus^C A$ denote the abelian group $I \oplus A$ with product defined by 2.2.1.

Proposition 2.3. *The natural isomorphism*

$$B \cong I \oplus A$$

of vector spaces is a unital ring isomorphism if and only if the following holds:

$$xC(y, z) - C(xy, z) + C(x, yz) - C(x, y)z = 0$$

for all $x, y, z \in A$.

Proof. We have defined two isomorphisms of vector spaces ϕ, ψ:

$$\phi(x) = (x - sp(x), p(x))$$

and

$$\psi(u, x) = u + s(x).$$

We define a product on the direct sum $I \oplus A$ using ϕ and ψ:

$$(u, x) \times (v, y) = \phi(\psi(u, x)\psi(v, y)) = \phi((u + s(x))(v + s(y))) =$$

$$\phi(uv+us(y)+s(x)v+s(x)s(y))=$$
$$(us(y)+s(x)v+s(x)s(y)-s(xy),xy)=(uy+xv+C(x,y),xy).$$

Here we define

$$uy=us(y)$$

and

$$xv=s(x)v.$$

One checks that

$$\phi(1)=(1-sp(1),1)=(0,1)=\mathbf{1}$$

and

$$\mathbf{1}(u,x)=(u,x)\mathbf{1}=(u,x)$$

for all $(u,x)\in I\oplus A$. It follows the morphism ϕ is unital. Since $C(x+y,z)=C(x,z)+C(y,z)$ and $C(x,y+z)=C(x,y)+C(x,z)$ the following holds:

$$(u,x)((v,y)+(w,z))=(u,x)(v,y)+(u,x)(w,z)$$

and

$$((v,y)+(w,z))(u,x)=(v,y)(u,x)+(w,z)(u,x).$$

Hence the multiplication is distributive over addition. Hence for an arbitrary section s of p of vector spaces mapping the identity to the identity it follows the multiplication defined above always has a left and right unit and is distributive. We check when the multiplication is associative.

$$((u,x)(v,y))(w,z)=(uyz+xvz+xyw+C(x,y)z+C(xy,z),xyz).$$

Also

$$(u,x)((v,y)(w,z))=(uyz+xvz+xyw+xC(y,z)+C(x,yz),xyz).$$

It follows the multiplication is associative if and only if the following equation holds for the element C:

$$xC(y,z)-C(xy,z)+C(x,yz)-C(x,y)z=0$$

for all $x,y,z\in A$. The claim follows. □

Let

$$xC(y,z)-C(xy,z)+C(x,yz)-C(x,y)z=0. \tag{2.3.1}$$

be the *cocycle condition.*

Definition 2.4. Let $\mathrm{exan}_k(A,I)$ be the set of elements $C\in\mathrm{Hom}_k(A\otimes_k A,I)$ satisfying the cocycle condition 2.3.1.

Proposition 2.5. *Equation 2.3.1 holds for all $x,y,z\in A$:*

Proof. We get:

$$xC(y,z)=s(x)s(y)s(z)-s(x)s(yz).$$
$$C(xy,z)=s(xy)s(z)-s(xyz).$$
$$C(x,yz)=s(x)s(yz)-s(xyz),$$

and

$$C(x,y)z=s(x)s(y)s(z)-s(xy)s(z).$$

We get

$$xC(y,z)-C(xy,z)+C(x,yz)-C(x,y)z=s(x)s(y)s(z)-s(x)s(yz)-$$

$$s(xy)s(z) + s(xyz) + s(x)s(yz) - s(xyz) - s(x)s(y)s(z) + s(xy)s(z) = 0$$

and the claim follows. □

Corollary 2.6. *The morphism $\phi : B \to I \oplus^C A$ is an isomorphism of unital associative k-algebras.*

Proof. This follows from Proposition 2.5 and Proposition 2.3. □

Hence there is always a commutative diagram of exact sequences

$$\begin{tikzcd}
0 \arrow[r] & I \arrow[r] \arrow[d, "="] & B \arrow[r] \arrow[d, "\cong"] & A \arrow[r] \arrow[d, "="] & 0 \\
0 \arrow[r] & I \arrow[r, "i"] & I \oplus^C A \arrow[r, "p"] & A \arrow[r] & 0
\end{tikzcd}$$

where the middle vertical morphism is an isomorphism associative unital k-algebras.

Define the following left and right A-action on the ideal I:

$$xu = s(x)u, ux = us(x)$$

where s is the section of p and $x \in A$, $u \in I$. Recall $I^2 = 0$.

Proposition 2.7. *The actions defined above give the ideal I a left and right A-module structure. The structure is independent of choice of section s.*

Proof. One checks that for any $x, y \in A$ and $u, v \in I$ the following holds:

$$(x + y)u = xu + yu, x(u + v) = xu + xv, 1u = 1.$$

Also

$$(xy)u - x(yu) = s(xy)u - s(x)s(y)u = (s(xy) - s(x)s(y))u = 0$$

since $I^2 = 0$. It follows $(xy)u = x(yu)$ hence I is a left A-module. A similar argument prove I is a right A-module. Assume t is another section of p. It follows

$$s(x)u - t(x)u = (s(x) - t(x))u = 0$$

since $I^2 = 0$. It follows $s(x)u = t(x)u$. Similarly $us(x) = ut(x)$ hence s and t induce the same structure of A-module on I and the Proposition is proved. □

We have proved the following Theorem: Let A be any associative unital k-algebra and let I be a left and right A-module. Let $C : A \otimes_k A \to I$ be a morphism satisfying the cocycle condition 2.3.1.

Theorem 2.8. *The exact sequence*

$$0 \to I \to I \oplus^C A \to A \to 0$$

is a square zero extension of A with the module I. Moreover any square zero extension of A with I arise this way for some morphism $C \in \mathrm{Hom}_k(A \otimes_k A, I)$ satisfying Equation 2.3.1.

Proof. The proof follows from the discussion above. □

Let

$$0 \to I \to E \to A \to 0$$

with $i : I \to E$ and $p : E \to A$ and

$$0 \to J \to F \to B \to 0$$

with $j : J \to F$ and $q : F \to B$ be square zero extensions of associative k-algebras A, B with left and right modules I, J. This means the sequences are exact and the following holds: $i(I)^2 = j(J)^2 = 0$. A triple (w, u, v) of maps of k-vector spaces giving rise to a commutative diagram of exact sequences

$$\begin{tikzcd}
0 \arrow[r] & I \arrow[r, "i"] \arrow[d, "w"] & E \arrow[r, "p"] \arrow[d, "u"] & A \arrow[r] \arrow[d, "v"] & 0 \\
0 \arrow[r] & J \arrow[r, "j"] & F \arrow[r, "q"] & B \arrow[r] & 0
\end{tikzcd}$$

is a morphism of extensions if u and v are maps of k-algebras and w is a map of left and right modules. This means

$$w(x + y) = w(x) + w(y), w(ax) = v(a)w(x), w(xa) = w(x)v(a)$$

for all $x, y \in I$ and $a \in A$.

We say two square zero extensions

$$0 \to I \to E \to A \to 0$$

and

$$0 \to I \to F \to A \to 0$$

are *equivalent* if there is an isomorphism $\phi : E \to F$ of k-algebras making all diagrams commute.

Definition 2.9. Let $\mathrm{Exan}_k(A, I)$ denote the set of all isomorphism classes of square zero extensions of A by I.

Theorem 2.10. *Let $C(A)$ be the center of A. The set $\mathrm{exan}_k(A, I)$ is a left and right module over $C(A)$. Moreover there is a bijection*

$$\mathrm{Exan}_k(A, I) \cong \mathrm{exan}_k(A, I)$$

of sets.

Proof. We first prove $\mathrm{exan}_k(A, I)$ is a left and right $C(A)$-module. Let $C, D \in \mathrm{exan}_k(A, I)$. This means $C, D \in \mathrm{Hom}_k(A \otimes_k A, I)$ are elements satisfying the cocycle condition 2.3.1. let $a, b \in C(A) \subseteq A$ be elements. Define aC, Ca as follows:

$$(aC)(x, y) = aC(x, y)$$

and

$$(Ca)(x, y) = C(x, y)a.$$

We see

$$\begin{gathered} x(aC)(y, x) - (aC)(xy, z) + (aC)(x, yz) - (aC)(x, y)z = \\ a(xC(y, z) - C(xy, z) + C(x, yz) - C(x, y)z) = a(0) = 0 \end{gathered}$$

hence $aC \in \mathrm{exan}_k(A, I)$. Similarly one proves $Ca \in \mathrm{exan}_k(A, I)$ hence we have defined a left and right action of $C(A)$ on the set $\mathrm{exan}_k(A, I)$. Given $C, D \in \mathrm{exan}_k(A, I)$ define

$$(C + D)(x, y) = C(x, y) + D(x, y).$$

One checks that $C + D \in \mathrm{exan}_k(A, I)$ hence $\mathrm{exan}_k(A, I)$ has an addition operation. One checks the following hold:

$$\begin{gathered} a(C + D) = aC + aD, (C + D)a = Ca + Da, \\ (a + b)C = aC + bC, C(a + b) = Ca + Cb, \end{gathered}$$

$$a(bC) = (ab)C, C(ab) = (Ca)b, 1C = C1 = C,$$

hence the set $\mathrm{exan}_k(A, I)$ is a left and right $C(A)$-module. Define the following map: Let $[B] = [I \oplus^C A] \in \mathrm{Exan}_k(A, I)$ be an equivalence class of a square zero extension. Define

$$\phi : \mathrm{Exan}_k(A, I) \to \mathrm{exan}_k(A, I)$$

by

$$\phi[B] = \phi[I \oplus^C A] = C.$$

We prove this gives a well defined map of sets: Assume $[I \oplus^C A]$ and $[I \oplus^D A]$ are two elements in $\mathrm{Exan}_k(A, I)$. Note: We use brackets to denote isomorphism classes of extensions. The two extensions are equivalent if and only if there is an isomorphism

$$f : I \oplus^C A \to I \oplus^D A$$

of k-algebras such that all diagrams are commutative. This means

$$f(u, x) = (u, x)$$

for all $(u, x) \in I \oplus^C A$. We get

$$f((u, x)(v, y)) = f(u, x)f(v, y).$$

This gives the equality

$$(uy + xv + C(x, y), xy) = (uy + xv + D(x, y), xy)$$

for all $(u, x), (v, y) \in I \oplus^C A$. Hence $\phi[I \oplus^C A] = C = D = \phi[I \oplus^D A]$ and the map ϕ is well defined. It is clearly an injective map. It is surjective by Theorem 2.8 and the claim of the Theorem follows. □

Theorem 2.10 shows there is a structure of left and right $C(A)$-module on the set of equivalence classes of extensions $\mathrm{Exan}_k(A, I)$. The structure as left $C(A)$-module agrees with the one defined in [3].

Let $\phi \in \mathrm{Hom}_k(A, I)$. Let $C^\phi \in \mathrm{Hom}_k(A \otimes_k A, I)$ be defined by

$$C^\phi(x, y) = x\phi(y) - \phi(xy) + \phi(x)y.$$

One checks that $C^\phi \in exan_k(A, I)$ for all $\phi \in \mathrm{Hom}_k(A, I)$.

Definition 2.11. Let $\mathrm{exan}_k^{inn}(A, I)$ be the subset of $\mathrm{exan}_k(A, I)$ of maps C^ϕ for $\phi \in \mathrm{Hom}_k(A, I)$.

Lemma 2.12. *The set* $\mathrm{exan}_k^{inn}(A, I) \subseteq \mathrm{exan}_k(A, I)$ *is a left and right sub* $C(A)$*-module.*

Proof. The proof is left to the reader as an exercise. □

Definition 2.13. Let $\mathrm{Exan}_k^{inn}(A, I) \subseteq \mathrm{Exan}_k(A, I)$ be the image of $\mathrm{exan}_k^{inn}(A, I)$ under the bijection $\mathrm{exan}_k(A, I) \cong \mathrm{Exan}_k(A, I)$.

It follows $\mathrm{Exan}_k^{inn}(A, I) \subseteq \mathrm{Exan}_k(A, I)$ is a left and right sub $C(A)$-module.

Recall the definition of the *Hochschild complex*:

Definition 2.14. Let A be an associative k-algebra and let I be a left and right A-module. Let $C^p(A, I) = \mathrm{Hom}_k(A^{\otimes p}, I)$. Let $d^p : C^p(A, I) \to C^{p+1}(A, I)$ be defined as follows:

$$d^p(\phi)(a_1 \otimes \cdots \otimes a_{p+1}) = a_1\phi(a_2 \otimes \cdots \otimes a_{p+1}) +$$

$$\sum_{1\leq i\leq p}(-1)^i\phi(a_1\otimes\cdots\otimes a_ia_{i+1}\otimes\cdots\otimes a_{p+1})+(-1)^{p+1}\phi(a_1\otimes\cdots\otimes a_p)a_{p+1}.$$

We let $\mathrm{HH}^i(A,I)$ denote the i'th cohomology of this complex. It is the *i'th Hochschild cohomology* of A with values in I.

Proposition 2.15. *There is an exact sequence*

$$0\rightarrow \mathrm{Exan}_k^{inn}(A,I)\rightarrow \mathrm{Exan}_k(A,I)\rightarrow \mathrm{HH}^2(A,I)\rightarrow 0$$

of left and right $C(A)$-modules.

Proof. The proof is left to the reader as an exercise. □

Example 2.16. *Characteristic classes of L-connections.*

Let A be a commutative k-algebra and let $\alpha : L\rightarrow \mathrm{Der}_k(A)$ be a Lie-Rinehart algebra. Let W be a left A-module with an L-connection $\nabla : L\rightarrow \mathrm{End}_k(W)$. In [6] we define a characteristic class $c_1(E)\in \mathrm{H}^2(L|_U,\mathcal{O}_U)$ when W is of finite presentation, $U\subseteq \mathrm{Spec}(A)$ is the open set where W is locally free and $\mathrm{H}^2(L|_U,\mathcal{O}_U)$ is the Lie-Rinehart cohomology of $L|_U$ with values in $\mathcal{O}_U$. If L is locally free it follows $\mathrm{H}^2(L,A)\cong \mathrm{Ext}^2_{U(L)}(A,A)$ where $U(L)$ is the generalized universal enveloping algebra of L. There is an obvious structure of left and right $U(L)$-module on $\mathrm{End}_k(A)$ and an isomorphism

$$\mathrm{HH}^2(U(L),\mathrm{End}_k(A))\cong \mathrm{Ext}^2_{U(L)}(A,A)$$

of abelian groups. The exact sequence 2.15 gives a sequence

$$0\rightarrow \mathrm{Exan}_k^{inn}(U(L),\mathrm{End}_k(A))\rightarrow \mathrm{Exan}_k(U(L),\mathrm{End}_k(A))\rightarrow$$
$$\mathrm{Ext}^2_{U(L)}(A,A)\rightarrow 0$$

with $A=U(L)$ and $I=\mathrm{End}_k(A)$. If we can construct a lifting

$$\tilde{c}_1(W)\in \mathrm{Exan}_k(U(L),\mathrm{End}_k(A))$$

of the class

$$c_1(W)\in \mathrm{Ext}^2_{U(L)}(A,A)=\mathrm{HH}^2(U(L),\mathrm{End}_k(A))$$

we get a generalization of the characteristic class from [6] to arbitrary Lie-Rinehart algebras L. This problem will be studied in a future paper on the subject (see [7]).

Example 2.17. *Non-commutative Kodaira-Spencer maps*

Let A be an associative k-algebra and let M be a left A-module. Let $D^1(A)\subseteq \mathrm{End}_k(A)$ be the *module of first order differential operators* on A. It is defined as follows: An element $\partial\in \mathrm{End}_k(A)$ is in $D^1(A)$ if and only if $[\partial,a]\in D^0(A)=A\subseteq \mathrm{End}_k(A)$ for all $a\in A$. Define the following map:

$$f: D^1(A)\rightarrow \mathrm{Hom}_k(A,\mathrm{End}_k(M))$$

by

$$f(\partial)(a,m)=[\partial,a]m=(\partial(a)-a\partial(1))m.$$

Here $\partial\in D^1(A), a\in A$ and $m\in M$. Since $[\partial,a]\in A$ we get a well defined map. Let for any $a\in A$ and $m\in M$ $\phi_a(m)=am$. It follows $\phi_a\in \mathrm{End}_k(M)$ is an endomorphism of M. We get

$$f(\partial)(ab,m)=(\partial(ab)-ab\partial(1))m=(\partial\phi_{ab}-\phi_{ab}\partial)(1)m=$$
$$(\partial\phi_{ab}-\phi_a\partial\phi_b+\phi_a\partial\phi_b-\phi_{ab}\partial)(1)m=$$

$$(\partial\phi_a - \phi_a\partial)\phi_b(1)m + \phi_a(\partial\phi_b - \phi_b\partial)(1)m =$$
$$f(\partial)(a, bm) + af(\partial)(b, m).$$

Hence

$$f(\partial)(ab) = af(\partial)(b) + f(\partial)(a)b$$

for all $\partial \in D^1(A)$ and $a, b \in A$. The Hochschild complex gives a map

$$d^1 : \mathrm{Hom}_k(A, \mathrm{End}_k(M)) \to \mathrm{Hom}_k(A \otimes A, \mathrm{End}_k(M))$$

and

$$ker(d^1) = \mathrm{Der}_k(A, \mathrm{End}_k(M)).$$

It follows we get a map

$$f : D^1(A) \to \mathrm{Der}_k(A, \mathrm{End}_k(M)).$$

We get an induced map

$$f : D^1(A) \to \mathrm{HH}^1(A, \mathrm{End}_k(M)) = \mathrm{Ext}^1_A(M, M).$$

Lemma 2.18. *The following holds:* $f(D^0(A)) = f(A) = 0$

Proof. The proof is left to the reader as an exercise. □

One checks that $D^1(A)/D^0(A) = D^1(A)/A \cong \mathrm{Der}_k(A)$. It follows we get an induced map

$$g : \mathrm{Der}_k(A) = D^1(A)/D^0(A) \to \mathrm{Ext}^1_A(M, M)$$

the *non-commutative Kodaira-Spencer map.*

Lemma 2.19. *Assume A is commutative. The following holds:*

(2.19.1) $\mathbb{V}_M = ker(g) \subseteq \mathrm{Der}_k(A)$ *is a Lie-Rinehart algebra.*

(2.19.2) $g(\delta) = 0 \iff \exists\phi \in \mathrm{End}_k(M), \phi(am) = a\phi(m) + \delta(a)m.$

(2.19.3) $\exists\nabla \in \mathrm{Hom}_k(\mathbb{V}_M, \mathrm{End}_k(M))$ *with* $\nabla(\delta)(am) = a\nabla(\delta)(m) + \delta(a)m.$

(2.19.4) $\mathbb{V}_M$ *is the maximal Lie-Rinehart algebra satisfying 2.19.3.*

Proof. We first prove 2.19.1: Assume $g(\delta) = g(\eta) = 0$. By definition this is if and only if there are maps $\phi, \psi \in \mathrm{End}_k(M)$ such that the following holds:

$$d^0\phi = g(\delta) \tag{2.19.5}$$

$$d^0\psi = g(\eta). \tag{2.19.6}$$

One checks that condition 2.19.5 and 2.19.6 hold if and only if the following hold:

$$\phi(am) = a\phi(m) + \delta(a)m$$

and

$$\psi(am) = a\psi(m) + \eta(a)m.$$

We claim : $d^0[\delta, \eta] = g([\delta, \eta])$: We get

$$[\phi, \psi](am) = \phi\psi(am) - \psi\phi(am) =$$
$$\phi(a\psi(m) + \eta(a)m) - \psi(a\phi(m) + \delta(a)m) =$$
$$a\phi\psi(m) + \delta(a)\psi(m) + \eta(a)\phi(m) + \delta\eta(a)m -$$
$$a\psi\phi(m) - \eta(a)\phi(m) - \delta(a)\psi(m) - \eta\delta(a)m =$$
$$a[\phi, \psi](m) + [\delta, \eta](a)m.$$

Hence $g([\delta, \eta]) = 0$ and $\mathbb{V}_M \subseteq \mathrm{Der}_k(A)$ is a k-Lie algebra. It is an A-module since g is A-linear, hence it is a Lie-Rinehart algebra. Claim 2.19.1 is proved. Claim 2.19.2

and 2.19.3 follows from the proof of 2.19.1. Claim 2.19.4 is obvious and the Lemma is proved. □

The Lie-Rinehart algebra $\mathbb{V}_M$ is the *linear Lie-Rinehart algebra* of M.

Let in the following E be a left and right A-module.

Definition 2.20. Let

$$\mathcal{J}^1_I(E) = I \otimes_A E \oplus E$$

be the *first order* I*-jet bundle of* E.

Pick a derivation $d \in \mathrm{Der}_k(A, I)$ of left and right modules. This means

$$d(xy) = xd(y) + d(x)y$$

for all $x, y \in A$. Let $B^C = I \oplus^C A$ and define the following left B^C-action on $\mathcal{J}^1_I(E)$:

$$(u, x)(w \otimes e, f) = (u \otimes f + xw \otimes e + d(x) \otimes f, xf)$$

for any elements $(u, x) \in B^C$ and $(w \otimes e, f) \in \mathcal{J}^1_I(E)$.

Proposition 2.21. *The abelian group* $\mathcal{J}^1_I(E)$ *is a left* B^C*-module if and only if* $C(y, x) \otimes f = 0$ *for all* $y, x \in A$ *and* $f \in E$.

Proof. One easily checks that for any $a, b \in B^C$ and $l, j \in \mathcal{J}^1_I(E)$ the following hold:

$$(a + b)i = ai + bi$$

$$a(i + j) = ai + aj.$$

Moreover

$$\mathbf{1}i = i.$$

It remains to check that $a(bi) = (ab)i$. Let $a = (v, y) \in B^C$ and $b = (u, x) \in B^C$. Let also $i = (w \otimes e, f) \in \mathcal{J}^1_I(E)$. We get

$$a(bi) = (v, y)((u, x)(w \otimes e, f)) = (vx \otimes f + yu \otimes f + yxw \otimes e + d(yx) \otimes f, yxf).$$

We also get

$$(ab)i = (vx \otimes f + yu \otimes f + yxw \otimes e + d(yx) \otimes f + C(y, x) \otimes f, yxf).$$

It follows that

$$(ab)i - a(bi) = 0$$

if and only if

$$C(y, x) \otimes f = 0,$$

and the claim of the Proposition follows. □

Note the abelian group $\mathcal{J}^1_I(E)$ is always a left A-module and there is an exact sequence of left A-modules

$$0 \to I \otimes E \to \mathcal{J}^1_I(E) \to E \to 0$$

defining a characteristic class

$$c_I(E) \in \mathrm{Ext}^1_A(E, E \otimes I).$$

The class $c_I(E)$ has the property that $c_I(E) = 0$ if and only if E has an I-connection:

$$\nabla : E \to I \otimes E$$

with

$$\nabla(xe) = x\nabla(e) + d(x) \otimes e.$$

Let $J \subseteq I \subseteq B^C$ be the smallest two sided ideal containing $Im(C)$ where $C : A \otimes_k A \rightarrow I$ is the cocycle defining B^C. Let $D^C = B^C/J$ and $I^C = I/J$. We get a square zero extension

$$0 \rightarrow I^C \rightarrow D^C \rightarrow A \rightarrow 0$$

of A by the square zero ideal I^C. It follows $D^C = I^C \oplus A$ as abelian group. Since $\overline{C(x,y)} = 0$ in I^C it follows D^C has a well defined associative multiplication defined by

$$(u,x)(v,y) = (uy + xv, xy).$$

Also D^C is the largest quotient of B^C such that the ring homomorphism $B^C \rightarrow D^C$ fits into a commutative diagram of square zero extensions

$$\begin{tikzcd}
0 \arrow[r] & I \arrow[r] \arrow[d] & B^C \arrow[r] \arrow[d] & A \arrow[r] \arrow[d, "="] & 0 \\
0 \arrow[r] & I^C \arrow[r] & D^C \arrow[r] & A \arrow[r] & 0.
\end{tikzcd}$$

Definition 2.22. Let

$$\mathcal{J}^1_{I^C}(E) = I^C \otimes E \oplus E$$

be the *first order I^C-jet bundle of E.*

Example 2.23. *First order commutative jets.*

Let $k \rightarrow A$ be a commutative k-algebra and let $I \subseteq A \otimes_k A$ be the ideal of the diagonal. Let $\mathcal{J}^1_A = A \otimes A/I^2$ and $\Omega^1_A = I/I^2$. We get an exact sequence of left A-modules

$$0 \rightarrow \Omega^1_A \rightarrow \mathcal{J}^1_A \rightarrow A \rightarrow 0. \tag{2.23.1}$$

It follows $\mathcal{J}^1_A \cong \Omega^1_A \oplus A$ with the following product:

$$(\omega, a)(\eta, b) = (\omega a + b\eta, ab)$$

hence the sequence 2.23.1 splits. Let $\mathcal{J}^1_A(E) = \Omega^1_A \otimes E \oplus E$ be the first order Ω^1_A-jet of E. We get an exact sequence of left A-modules

$$0 \rightarrow \Omega^1_A \otimes E \rightarrow \mathcal{J}^1_A(E) \rightarrow E \rightarrow 0.$$

Since the sequence 2.23.1 splits it follows $\mathcal{J}^1_A(E)$ is a lifting of E to the first order jet $\mathcal{J}^1_A$.

3. Atiyah classes and Kodaira-Spencer classes

In this section we define and prove some properties of Atiyah classes and Kodaira-Spencer classes.

Let X be any scheme defined over an arbitrary basefield F and let $\operatorname{Pic}(X)$ be the *Picard group* of X. Let $\mathcal{O}^* \subseteq \mathcal{O}_X$ be the following subsheaf of abelian groups: For any open set $U \subseteq X$ the group $\mathcal{O}(U)^*$ is the multiplicative group of units in $\mathcal{O}_X(U)$. Define for any open set $U \subseteq X$ the following morphism:

$$\operatorname{dlog} : \mathcal{O}(U)^* \rightarrow \Omega^1_X(U)$$

defined by

$$\operatorname{dlog}(x) = d(x)/x,$$

where d is the universal derivation and $x \in \mathcal{O}(U)^*$.

Lemma 3.1. *The following hold:*

$$\mathrm{dlog}(xy) = \mathrm{dlog}(x) + \mathrm{dlog}(y)$$

for $x, y \in \mathcal{O}(U)^*$

Proof. The proof is left to the reader as an exercise. $\square$

Hence $\mathrm{dlog} : \mathcal{O}^* \to \Omega^1_X$ defines a map of sheaves of abelian groups. The map dlog induce a map on cohomology

$$\mathrm{dlog} : \mathrm{Pic}(X) = \mathrm{H}^1(X, \mathcal{O}^*) \to \mathrm{H}^1(X, \Omega^1_X)$$

and by definition

$$c_l(\mathcal{L}) = \mathrm{dlog}(\mathcal{L}) \in \mathrm{H}^1(X, \Omega^1_X).$$

Let $\mathcal{I} \subseteq \Omega^1_X$ be any sub $\mathcal{O}_X$-module and let $\mathcal{F} = \Omega^1_X/\mathcal{I}$ be the quotient sheaf. We get a derivation

$$d : \mathcal{O}_X \to \mathcal{F}$$

by composing with the universal derivation. We get a canonical map

$$\mathrm{H}^1(X, \Omega^1_X) \to \mathrm{H}^1(X, \mathcal{F})$$

and we let

$$\overline{c}_1(\mathcal{L}) \in \mathrm{H}^1(X, \mathcal{F})$$

be the image of $c_1(\mathcal{L})$ under this map.

Definition 3.2. The class $c_1(\mathcal{L}) \in \mathrm{H}^1(X, \Omega^1_X)$ is the *first Chern class* of the line bundle $\mathcal{L} \in \mathrm{Pic}(X)$. The class $\overline{c}_1(\mathcal{L}) \in \mathrm{H}^1(X, \mathcal{F})$ is the *generalized first Chern class* of $\mathcal{L}$.

Let $\mathcal{E}$ be any $\mathcal{O}_X$-module and consider the following sequence of sheaves of abelian groups:

$$0 \to \mathcal{F} \otimes \mathcal{E} \to \mathcal{J}^1_{\mathcal{F}}(\mathcal{E}) \to \mathcal{E} \to 0$$

where

$$\mathcal{J}^1_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \otimes \mathcal{E} \oplus \mathcal{E}$$

as sheaf of abelian groups. Let s be a local section of $\mathcal{O}_X$ and let $(x \otimes e, f)$ be a local section of $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E})$ over some open set U. Make the following definition:

$$s(x \otimes e, f) = (sx \otimes e + ds \otimes f, sf).$$

It follows the sequence

$$0 \to \mathcal{F} \otimes \mathcal{E} \to \mathcal{J}^1_{\mathcal{F}}(\mathcal{E}) \to \mathcal{E} \to 0$$

is a short exact sequence of sheaves of abelian groups. It is called the *Atiyah-Karoubi sequence.*

Definition 3.3. An $\mathcal{F}$-connection ∇ is a map

$$\nabla : \mathcal{E} \to \mathcal{F} \otimes \mathcal{E}$$

of sheaves of abelian groups with

$$\nabla(se) = s\nabla(e) + d(s) \otimes e.$$

Proposition 3.4. *The Atiyah-Karoubi sequence is an exact sequence of left* $\mathcal{O}_X$*-modules. It is left split by an* $\mathcal{F}$*-connection.*

Proof. We first show it is an exact sequence of left $\mathcal{O}_X$-modules. The $\mathcal{O}_X$-module structure is twisted by the derivation d, hence we must verify that this gives a well defined left $\mathcal{O}_X$-structure on $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E})$. Let $\omega = (x \otimes e, f)$ be a local section of $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E})$ and let s, t be local sections of $\mathcal{O}_X$. We get the following calculation:

$$(st)\omega = (st)(x \otimes e, f) = ((st)x \otimes e + d(st) \otimes f, (st)f) =$$

$$(stx \otimes e + sdt \otimes f + (ds)t \otimes f, stf) = (s(tx \otimes e + dt \otimes f) + ds \otimes tf, s(tf)) =$$

$$s(tx \otimes e + dt \otimes f, tf) = s(t(x \otimes e, f)) = s(t\omega).$$

It follows $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E})$ is a left $\mathcal{O}_X$-module and the sequence is left exact. Assume

$$s : \mathcal{E} \to \mathcal{J}_{\mathcal{F}}(\mathcal{E}) = \mathcal{F} \otimes \mathcal{E} \oplus \mathcal{E}$$

is a left splitting. It follows $s(e) = (\nabla(e), e)$ for e a local section of $\mathcal{E}$. It follows ∇ is a generalized connection and the Theorem is proved. □

Note: If $\mathcal{I} = 0$ we get $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E}) = \mathcal{J}^1_X(\mathcal{E})$ is the first order jet bundle of $\mathcal{E}$ and the exact sequence above specialize to the well known *Atiyah sequence*:

$$0 \to \Omega^1_X \otimes \mathcal{E} \to \mathcal{J}^1_X(\mathcal{E}) \to \mathcal{E} \to 0.$$

The Atiyah sequence is left split by a connection

$$\nabla : \mathcal{E} \to \Omega^1_X \otimes \mathcal{E}.$$

The $\mathcal{O}_X$-module $\mathcal{J}^1_{\mathcal{F}}(\mathcal{E})$ is the *generalized first order jet bundle of* $\mathcal{E}$.

Definition 3.5. The characteristic class

$$\mathrm{AT}(\mathcal{E}) \in \mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{E}, \mathcal{F} \otimes \mathcal{E})$$

is called the *Atiyah class* of $\mathcal{E}$.

The class $\mathrm{AT}(\mathcal{E})$ is defined for an arbitrary $\mathcal{O}_X$-module $\mathcal{E}$ and an arbitrary sub module $\mathcal{I} \subseteq \Omega^1_X$.

Assume $\mathcal{E} = \mathcal{L} \in \mathrm{Pic}(X)$ is a line bundle on X. We get isomorphisms

$$\mathrm{Ext}_{\mathcal{O}_X}(\mathcal{L}, \mathcal{L} \otimes \mathcal{F}) \cong \mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{L}^* \otimes \mathcal{L} \otimes \mathcal{F}) \cong$$

$$\mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \mathcal{F}) \to \mathrm{H}^1(X, \mathcal{F}).$$

We get a morphism

$$\phi : \mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{L}, \mathcal{L} \otimes \mathcal{F}) \to \mathrm{H}^1(X, \mathcal{F}).$$

Proposition 3.6. *The following hold:*

$$\phi(\mathrm{AT}(\mathcal{L})) = \overline{c}_1(\mathcal{L}).$$

Hence the Atiyah class calculates the generalized first Chern class of a line bundle.

Proof. Let $\mathcal{I} = 0$. It is well known that $\mathrm{AT}(\mathcal{L})$ calculates the first Chern class $c_1(\mathcal{L})$. From this the claim of the Proposition follows. □

Let T_X be the tangent sheaf of X. It has the property that for any open affine set $U = \mathrm{Spec}(A) \subseteq X$ the local sections $T_X(U)$ equals the module $\mathrm{Der}_F(A)$ of derivations of A. Let $\mathbb{V}_{\mathcal{E}} \subseteq T_X$ be the subsheaf of local sections ∂ of T_X with the following property: The section $\partial \in T_X(U)$ lifts to a local section $\nabla(\partial)$ of $\mathrm{End}_F(\mathcal{E}|_U)$ with the following property:

$$\nabla(\partial) : \mathcal{E}|_U \to \mathcal{E}|_U$$

satisfies

$$\nabla(\partial)(se) = s\nabla(\partial)(e) + \partial(s)e.$$

It follows $\mathbb{V}_{\mathcal{E}} \subseteq T_X$ is a sheaf of Lie-Rinehart algebras - the *Kodaira-Spencer sheaf* of $\mathcal{E}$.

Define for any local sections a, b of $\mathcal{O}_X$,∂ of $\mathbb{V}_{\mathcal{E}}$ and e of $\mathcal{E}$ the following:

$$L(a,\partial)(e) = a\nabla(\partial)(e) - \nabla(a\partial)(e).$$

Lemma 3.7. *It follows* $L(a,\partial) \in \operatorname{End}_{\mathcal{O}_U}(\mathcal{E}|_U)$.

Proof. The following hold:

$$L(a,\partial)(be) = a\nabla(\partial)(be) - \nabla(a\partial)(be) =$$

$$a(b\nabla(\partial)(e) + \partial(b)e) - b\nabla(a\partial)(e) - a\partial(b)e =$$

$$ab\nabla(\partial)(e) + a\partial(b)e - b\nabla(a\partial)(e) - a\partial(b)e =$$

$$b(a\nabla(\partial)(e) - \nabla(a\partial)(e)) = b(a\nabla(\partial) - \nabla(a\partial))(e) =$$

$$bL(a,\partial)(e)$$

and the Lemma is proved. □

Lemma 3.8. *The following formula hold:*

$$L(ab,\partial) = aL(b,\partial) + L(a,b\partial)$$

for all local sections a, b *and* ∂.

Proof. We get

$$L(ab,\partial) = ab\nabla(\partial) - \nabla(ab\partial) =$$

$$ab\nabla(\partial) - a\nabla(b\partial) + a\nabla(b\partial) - \nabla(ab\partial) =$$

$$a(b\nabla(\partial) - \nabla(b\partial)) + (a\nabla - \nabla a)(b\partial) =$$

$$aL(b,\partial) + L(a,b\partial),$$

and the Lemma is proved. □

Let $\operatorname{LR}(\mathbb{V}_{\mathcal{E}}) = \operatorname{End}_{\mathcal{O}_X}(\mathcal{E}) \oplus \mathbb{V}_{\mathcal{E}}$ be the *linear Lie-Rinehart algebra* of $\mathcal{E}$. Let $\operatorname{LR}(\mathbb{V}_{\mathcal{E}})$ have the following left $\mathcal{O}_X$-module structure:

$$a(\phi,\partial) = (a\phi + L(a,\partial), a\partial).$$

Here a, ϕ and ∂ are local sections of $\mathcal{O}_X$, $\operatorname{End}_{\mathcal{O}_X}(\mathcal{E})$ and $\mathbb{V}_{\mathcal{E}}$. We twist the trivial $\mathcal{O}_X$ structure on $\operatorname{End}_{\mathcal{O}_X}(\mathcal{E}) \oplus \mathbb{V}_{\mathcal{E}}$ with the element L. We get a sequence of sheaves of abelian groups

$$0 \rightarrow \operatorname{End}_{\mathcal{O}_X}(\mathcal{E}) \rightarrow^i \operatorname{LR}(\mathbb{V}_{\mathcal{E}}) \rightarrow^p \mathbb{V}_{\mathcal{E}} \rightarrow 0$$

where i and p are the canonical maps. An $\mathcal{O}_X$-linear map

$$\nabla : \mathbb{V}_{\mathcal{E}} \rightarrow \operatorname{End}_F(\mathcal{E})$$

satisfying

$$\nabla(\partial)(ae) = a\nabla(\partial)(e) + \partial(a)e$$

is a $\mathbb{V}_{\mathcal{E}}$-*connection* on $\mathcal{E}$.

Proposition 3.9. *The sequence defined above is an exact sequence of left* $\mathcal{O}_X$*-modules. It is left split by a* $\mathbb{V}_{\mathcal{E}}$*-connection* ∇.

Proof. We need to check that $\mathrm{LR}(\mathbb{V}_{\mathcal{E}})$ has a well defined left $\mathcal{O}_X$-module structure. By definition

$$a(\phi, \partial) = (a\phi + L(a, \partial), a\partial).$$

We get

$$(ab)x = (ab)(\phi, \partial) = ((ab)\phi + L(ab, \partial), (ab)\partial) =$$
$$(ab\phi + aL(b, \partial) + L(a, b\partial), ab\partial) =$$
$$a(b\phi + L(b, \partial), b\partial) = a(b(\phi, \partial)) = a(bx)$$

and it follows the sequence is a left exact sequence of $\mathcal{O}_X$-modules. If

$$s : \mathbb{V}_{\mathcal{E}} \to \mathrm{End}_{\mathcal{O}_X}(\mathcal{E}) \oplus \mathbb{V}_{\mathcal{E}} = \mathrm{LR}(\mathbb{V}_{\mathcal{E}})$$

is a section it follows $s(e) = (\nabla(e), e)$. One checks that ∇ is a $\mathbb{V}_{\mathcal{E}}$-connection, and the Theorem is proved. □

Definition 3.10. We get a characteristic class

$$\mathrm{KS}(\mathcal{E}) \in \mathrm{Ext}^1_{\mathcal{O}_X}(\mathbb{V}_{\mathcal{E}}, \mathrm{End}_{\mathcal{O}_X}(\mathcal{E}))$$

the *Kodaira-Spencer class* of $\mathcal{E}$.

Assume $\mathbb{V}_{\mathcal{E}}$ is locally free and $\mathcal{E} = \mathcal{L} \in \mathrm{Pic}(X)$ is a line bundle on X. Assume also $\mathbb{V}^*_{\mathcal{E}} = \mathcal{F} = \Omega^1_X/\mathcal{I}$ for some submodule $\mathcal{I}$. We get the following calculation:

$$\mathrm{Ext}^1_{\mathcal{O}_X}(\mathbb{V}_{\mathcal{E}}, \mathrm{End}_{\mathcal{O}_X}(\mathcal{L})) \cong \mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \mathrm{End}_{\mathcal{O}_X}(\mathcal{L}) \otimes \mathbb{V}^*_{\mathcal{E}}) \cong$$
$$\mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{O}_X, \mathrm{End}_{\mathcal{O}_X}(\mathcal{L}) \otimes \mathcal{F}) \to \mathrm{H}^1(X, \mathcal{F}).$$

We get a map

$$\psi : \mathrm{Ext}^1_{\mathcal{O}_X}(\mathbb{V}_{\mathcal{E}}, \mathrm{End}_{\mathcal{O}_X}(\mathcal{L})) \to \mathrm{H}^1(X, \mathcal{F})$$

of sheaves.

Proposition 3.11. *The following hold: There is an equality*

$$\psi(\mathrm{KS}(\mathcal{L})) = \overline{c}_1(\mathcal{L})$$

in $\mathrm{H}^1(X, \mathcal{F})$*. Hence the Kodaira-Spencer class calculates the class* $\overline{c}_1(\mathcal{L})$*.*

Proof. The proof is left to the reader as an exercise. □

We get the following diagram expressing the relationship between the characteristic classes defined above:

$$\begin{tikzcd}
\mathrm{Ext}^1_{\mathcal{O}_X}(\mathbb{V}_{\mathcal{L}}, \mathrm{End}_{\mathcal{O}_X}(\mathcal{L})) \arrow[dr, "\psi"] & & \\
& \mathrm{H}^1(X, \mathcal{F}) & \mathrm{Pic}(X) \arrow[l, "\overline{c}_1(-)"'] \\
\mathrm{Ext}^1_{\mathcal{O}_X}(\mathcal{L}, \mathcal{F} \otimes \mathcal{L}) \arrow[ur, "\phi"] & &
\end{tikzcd}$$

The following equation holds in $\mathrm{H}^1(X, \mathcal{F})$:

$$\phi(\mathrm{AT}(\mathcal{L})) = \psi(\mathrm{KS}(\mathcal{L})) = \overline{c}_1(\mathcal{L}).$$

References

[1] M. Andre, Homologie des algebres commutatives, *Grundlehren Math. Wiss.* no. 206 (1974)
[2] M. Atiyah, Complex analytic connections in fibre bundles, *Trans. AMS* no. 85 (1957)
[3] A. Grothendieck, EGA IV Etude locale de schemas et des morphismes de schemas, *Publ. Math. IHES* no. 20 (1964)
[4] M. Karoubi, Homologie cyclique et K-theorie, *Asterisque* no 149 (1987)
[5] H. Maakestad, A note on the principal parts on projective space and linear representations, *Proc. of the AMS* Vol. 133 no. 2 (2004)
[6] H. Maakestad, Chern classes and Lie-Rinehart algebras, *Indagationes Math.* (2008)
[7] H. Maakestad, Chern classes and Exan functors, *In progress* (2009)
[8] H. Maakestad, Principal parts on the projective line over arbitrary rings, *Manuscripta Math.* 126, no. 4 (2008)

Oslo, Norway
E-mail address: `h_maakestad@hotmail.com`